# Analytical Study on Motion Behavior of Non-Spherical Particles in Incompressible Fluid with Presence of Electrostatic Force

A. R. Noorpoor<sup>1</sup>
A. R. Noorpoor<sup>1</sup>
I University of Tehran
Received: 6 April 2015 Accepted: 30 April 2015 Published: 15 May 2015

#### 7 Abstract

8 In this paper, the accelerated motion of non-spherical particles in an incompressible fluid in

<sup>9</sup> both the presence and the absence of electrostatic force was investigated. Differential

<sup>10</sup> transformation method (DTM) and a FORTRAN code was used to calculate the instantaneous

velocity of particles. Regarding particles? instantaneous velocity in the absence of electrostatic

<sup>12</sup> force, DTM approach was resulted in a proper accordance with previous studies which utilized

<sup>13</sup> variational iteration method (VIM). In addition, a good agreement between DTM and VMT

<sup>14</sup> was seen as sphericity of particles was varied from 0.5 to 0.9. The results showed that falling

velocity increased with increasing sphericity. Moreover, the presence of electrostatic force (by
 assuming the electrical load equal to 1 micro colon) was compared to the one with no

<sup>17</sup> electrostatic force. The results showed that the falling velocity was decreased by 23.33

18

19 Index terms— electrostatic, particles, acceleration motion, fluid, liquid.

# 20 1 Introduction

where?? ?? is particle mass,??is velocity, ??is time, ?? is original analytical function, and?? ?? is the resultant of forces that are applied to the particle, including gravity, buoyancy, drag, virtual mass force, the Basset force, and lift force.

Equation (1) has been solved numerically in different studies by various methods, for example the finite difference method [1]. Some analytical methods which were applied for analysis of the acceleration of spherical and non-spherical particles motion in Newtonian fluids was addressed in [2][3][4][5][6][7][8][9][10]. Jalaal and Ganji [3]studied spherical and non-spherical particles motion in unsteady state at Newtonian media. They used friction coefficient governed by Chhabra and Ferreira's equations [11]for a range of Reynolds numbers by using the

homotopy perturbation method (HPM). Jalaal et al. [6]studied non-spherical particles motion at the Newtonian
media using the VIM and friction equations derived by Chien [12]. In another researches, Jalaal et al. [4] and
Jalaal and Ganji [5]studied non-spherical particles by using the HPM.

Stokes [13]assigned following equation for drag coefficient of a sedimenting particle. The equation is derived

40 for a flow field that is totally dominated by viscous diffusion as belowC D = 24 Re (2)

where C D is drag coefficient and Re is Reynolds number. This equation denied the effect of inertia and is accepted for???? < 0.4. Therefore, Oseen [14], assuming the effect of inertia, completed the Stokes's equation as follow?? ?? = 24 ???? ?1 + 3 16 ?????(3) Most of previous researches on spherical and non-spherical particles carried out experimentally and only a few of them were analytically investigated the solution of motion equations. In this regard, Proudman and Pearson [15]proposed????  $2 \times \ln$  (????) parameter to consider the behavior of drag coefficient for spherical particles and then Sano [16]completed aforementioned equation. Lovalenti and Brady [17] used the Kim and Karilla's equations [18]to solve directly the behavior of applied forces on particles and derived different parameters of time reduction, Including?? ?1, ?? ?5/2,?? ??? , ?? ?2. ?? ??? , ?? ?1/2, and ?? ?2, depending on initial conditions. Ferreira and Chhabra [11]achieved the following equation for 0 ? Re ?10 5

51 **2 S** 

# <sup>52</sup> 3 Global Journal of Researches in Engineering

53 ?? ?? = 24 ???? ?1 + 1 48 ?????(4)

Equation (4) showed a suitable accordance with results of experimental researches. Also, in many cases, a linear equation can describe the drag force very well. In some cases, Reynolds number has an average value and the liner and exponential terms are used in [11,12,[19] ??20] ??21] ??22].

64 Where ??, ??, and ?? are constants and depend on physical condition of the system.

<sup>65</sup> Different methods were proposed in previous studies to solve equation (6). Among those, VIM, a technique <sup>66</sup> based on repeated integration proposed by He ??24], was used successfully to solve linear and nonlinear equations <sup>67</sup> by different authors. This method is based on the true function by a general Lagrangian multiplier. DTM is <sup>68</sup> another method to solve the linear and non-linear terms proposed by Yaghoobi and Torabi [8]. Noorpoor and

<sup>69</sup> Nazari [25] used DTM to calculate the falling velocity of a particle in the acoustic field.

# 70 **4** II.

# 71 5 Materials and Methods

In this paper, equation (??) was solved by using the Yaghoobi and Torabi's equations [8]. The solving procedure
 included applying DTM by using a FORTRAN code.

By combining equation (??), Reynolds number (???? = ?????? ?? , ?? is dynamic viscosity), and drag coefficient equation, the final equation of a falling acceleration of non-spherical particle motion in an incompressible fluid was written as follow??? + 1 12 ???? 3 ??? ???? + 3.75??????? + 67.289?? 77 (?5.03?) 8 ???? 2 ???? 2 ? ???? ?1 ? ?? ?? ?? = 0(7)

where ? is sphericity of particle. For simplicity, this equation's coefficients were assigned as a, b, c and d in further equations. First term of equation (7) is to describe the added mass to a falling non-spherical particle in an incompressible fluid. The second and third term show the resistant force of particle's linear and non-linear motion respectively, and the fourth shows the applied gravity and buoyancy forces on the falling nonspherical particle in an incompressible fluid.?? = (?? + 1 12 ???? 3 ??)(8)?? = 3.75?????(9)?? = 67.289?? (?5.03?) 8 ???? 2 ?? (10) ?? = ???? ?1 ? ?? ?? ?(11)

So, equation (7) was rewritten as follow?? ???? ???? + ???? + ???? 2??? = 0, ??(0) = 0(12)

According to DTM conversion functions, equation (12) is provided as following transformed equation??(?? + 1)??(?? + 1) + ????(??) + ???? ??(??)??(?? ? 1) ?? ??=0 ? ? ?? × ??(??) = 0(13)

The value of ??(0) is equal to zero. Other value of ??(??) for k=1, 2, 3 ? are governed by above equation and can be calculated as bellow.??(1) = ?? ?? (14) ??(2) = ? 1 2 × ???? ?? 2 (15) ??(3) = 1 3! × ??(?? 2 ?2???? ) ?? 3 (16) ??(4) = ? 1 4! × ???? (?? 2 ?8???? ) ?? 4(17)??(5) = 1 5! × ??(?? 4 ?22?? 2 ???? +16?? 2 ?? 2 ) ?? 5 (18) ??(6) = ? 1 6! × ???? (?? 4 ?52?? 2 ???? +136?? 2 ?? 2 ) ?? 6(19)??(7) = 1 7! × ??(?? 6 ?114?? 4 ???? +720?? 2 ?? 2 ?? 2 ??2???? 3 ?? 3 ) ?? 7 (20) ??(8) = ? 1 8! × ???? (?? 6 ?240?? 4 ???? +3072?? 2 ?? 2 ?? 2 ?3968?? 3 ?? 3 ) ?? 8(21)

To calculate the velocity at each moment following equation is using?? ?? (??) = ? ? ?? ?? ?? ?? ?? ?? ?? ?? 94 (??), 0 ? ?? ? ?? ??

spherical particles in an incompressible fluid is derived as?? ?? (??) = ??(1)?? + ??(2)?? 2 + ??(3)?? 3 + ??(4)?? 4 + ?(23)

In order to study the impact of electrostatic force, Coulomb's lawis used as follow?? = ??  $2 4 \times$ ??  $2 \times$ ??  $0 \times$ ?? 98 2, ??  $0 = 8.85 \times 10$  ?12 (24)

Where ?? is electrostatic force, ?? is the signed magnitude of the charges, ?? is the distance between the charges, and ?? 0 is vacuum permittivity coefficient.m ???? ???? = ???? ?1 ? ?? ?? ?? ? 1 8 ???? 2 ???? ?? ?? 2 ? 1 12 ???? 3 ?? ???? ???? + ?? 2  $4 \times ?? \times ??$  0  $\times ??$  2 (25) Assuming that all varieties of equation (24) are independent of time, following equation can be written by rewriting terms of equation (??5) and inserting equation (7) in (11), and a non-linear deferential equation is formed:?? ???? ???? + ???? + ???? 2 ? ?? ?? = 0, ??(0) = 0 (26)

By applying the DTM one quation (26) the following equation is derived.

#### 

By considering equations numbers from (13) to (27),the term (d+e) was replaced to (d) to calculate U(1) to U (8). As a result:??(1) = (??+??) ?? (28) ??(2) = ? 1 2 × ??(??+??) ?? 2 (29) ??(3) = 1 3! × (??+??)(?? 2 ???(??+??)) ?? 3 (30) ??(4) = ? 1 4! × ??(??+??)(?? 2 ?8??(??+??)) ?? 4 (31) ??(5) = 1 5! × (??+??)(?? 4 ?2?? 2 ??(??+??)+16?? 2 (??+??) 2 ) ?? 5 (32) ??(6) = ? 1 6! × ??(??+??)(?? 4 ?52?? 2 ??(??+??)+136?? 2 (??+??) 2 ) ?? 6 (33) ??(7) = 1 7! × (??+??)(?? 6 ?114?? 4 ??(??+??)+720?? 2 ?? 2 (??+??) 2 ?272?? 3 13 (??+??) 3 ) ?? 7 (34) ??(8) = ? 1 8! × ??(??+??)(?? 6 ?240?? 4 ??(??+??)+3072?? 2 ?? 2 (??+??) 2 ?3968?? 14 3 (??+??) 3 ) ?? 8(35)

Finally, using final derived equations described above, the equation (23)was solved as the general equation using a FORTRAN code.

# 117 **7 III.**

#### <sup>118</sup> 8 Results and Discussion

The FORTRAN code is used for calculating instantaneous velocity,?? ?? (??), by assuming sphericity, density and equivalent diameterof particle equal to 0.9, 2100 ???? ?? 3 and 3mm respectively.

Figure ?? illustrated the changes in the instantaneous velocity for sphericity of 0.9 when the fluid was water 121 and no electrical load was applied. The falling velocity increased with increasing time. This increasing approach 122 followed the linear behavior at the beginning, until the velocity passed 0.1 m/sec point. However, as it can be 123 seen, the value of velocity reached to a constant value of 0.16 m/sec after 0.07 sec. Figure ?? Figure 2 and 3 124 compares instantaneous velocities in water and ethylene glycol as their densities differ. The same approach can 125 be seen regarding the increasing of velocity with time. However, the rate of increase in values of velocity was 126 higher for water as its density is lower than that of ethylene glycol. In addition, the rate in ethylene glycol's 127 representative diagram was increased in nonlinear approach in contrast with the one of water's. 128

Figure 4 represents the changes of particle's velocity when its sphericity changes. As it can be seen, the increase 129 in sphericity resulted in an increase in falling velocity. The trend of increasing velocity was approximately the 130 same before the velocity hit 0.04 m/sec independent of particles' sphericity. After this velocity, however, this 131 rate was higher for particles with higher sphericity. Figure 4 Falling velocity of particles with lower sphericity 132 reached a constant value in shorter times comparing the ones with lower sphericity. These results are consistent 133 with Yaghoobi's and Torabi's [8] and Jalaal's and Ganji's [3]. An error analyzing was carried out in order to 134 compare the results obtained in this study and Yaghoobi and Torabi's [8]; for particles with sphericity of 0.5 and 135 in absence of electrostatic force assuming the water as the incompressible fluid used for both studies, the value of 136 falling velocity differ 6.7%. Table 1 lists important points which can be taken form Figure 4 for falling velocities 137 of particles with different sphericities and the trend they increased. 138

#### <sup>139</sup> 9 Table 1

In order to investigate the effects an electrostatic force on particles' sedimentation By inserting the above equation 140 in equation (??), the equation (??5) is derived. micro colon applied to a non-spherical particle with sphericity 141 of 0.9 in water. The effects of electrostatic force in the motion behavior of particles in different distances between 142 electric charges (0.5 m and 1.0 m) are shows in Figure 5. It can be found from this figure that applying 143 electrostatic force reduced sedimentation velocity by 23.33% due to the formation of a resistant force between 144 particles. This reduction was more pronounced for shorter distance between the charges compared to the one 145 with longer distance. Also, it can be seen from Figure 5 that the effect of electrostatic force was approximately 146 constant with time especially after 0.04 sec. 147

148 IV.

## 149 **10** Conclusion

150 The differential transformation method (DTM) was used for analytical investigation of non-spherical particles' falling velocity in an incompressible fluid in both the presence and the absence of electrostatic force. Results 151 152 of this study indicated that, in general, the increasing curve of the sedimentation of non-spherical particles in 153 incompressible fluids mostly behave nonlinear until it reaches the constant value. In addition, the effects of changing fluid's density in sedimentation behavior of particles was investigated. The results showed the falling 154 velocity in water (the fluid with lower density) was higher than the one for ethylene glycol (the fluid with lower 155 density). Moreover, when the sphericity increased, the sedimentation rate increased even though this rate differed 156 for each sphericity. As results showed, particles with sphericity of 0.9 experienced the highest falling velocity and 157 increasing rate followed by those with sphericity of 0.7 and 0.5 respectively. After applying an electric load of 158

- 159 1 micro colon in with distant between charges of 0.5 m and 1.0 m, the falling velocity decreased as a resistant
- force was created. This decrease was more noticeable for the distance of 0.5 with decreasing about 23% in value
- 161 of falling velocity.

# 162 11 Global

## 163 **12 Tables**

Figure (1) -The changes of instantaneous velocity for sphericity of 0.9 with time in water and in absence of electrostatic force.  $^{1}$ 



Figure 1: Figure 2 Figure 3

#### 1

Sphericity Time of linear velocity Maximum linear velocity Start of constant Instantaneous velocity velocity m/sec m/sec sec sec ?=0.50.010.0306 0.030.06 ?= 0.70.0790.096 0.030.04?=0.90.06 0.1460.08 0.16

[Note: FiguresGlobal Journals Inc. (US) Guidelines Handbook 2015 www.GlobalJournals.org]

Figure 2: Table 1 :

165

 $<sup>^{1}</sup>$ © 2015 Global Journals Inc. (US)

- [Ganji ()] 'A semi-Analytical technique for nonlinear settling particle equation of Motion'. D D Ganji . Journal
   of Hydro-environment Research 2012. 6 p. .
- [Jalaal et al. ()] 'A series exact solution for one-dimensional non-linear particle equation of motion'. M Jalaal ,
   H Bararnia , G Domairry . *Powder Technology* 2011. 207 p. .
- [Ferreira and Chhabra] Accelerating motion of a vertically falling sphere in incompressible, J M Ferreira, R P
   Chhabra .
- [Chhabra and Ferreira ()] 'An analytical study of the motion of a sphere rolling down a smooth inclined plane
   in an incompressible Newtonian fluid'. R P Chhabra , J M Ferreira . *Powder technology* 1999. 104 p. .
- IZA [Jalaal and Ganji ()] 'An analytical study on motion of a sphere rolling down an inclined plane submerged in a
   Newtonian fluid'. M Jalaal , D D Ganji . *Powder Technology* 2010. 198 p. .
- IJalaal et al. ()] 'An analytical study on settling of non-spherical particles'. M Jalaal , D D Ganji , G Ahmadi .
   Asia-Pacific Journal of Chemical Engineering 2012. 7 p. .
- IJalaal et al. ()] 'Analytical investigation on acceleration motion of a vertically falling spherical particle in
   incompressible Newtonian media'. M Jalaal , D D Ganji , G Ahmadi . Advanced Powder Technology 2010. 21
   p. .
- [Yaghoobi and Torabi ()] 'Analytical solution for settling of non-spherical particles in incompressible Newtonian
   media'. H Yaghoobi , M Torabi . *Powder Technology* 2012. p. .
- [Proudman and Pearson ()] 'Expansions at small Reynolds numbers for the flow past a sphere and a circular
   cylinder'. I Proudman, J R A Pearson. Journal of Fluid Mechanics 1957. 2 p. .
- [Jalaal et al. ()] 'Homotopy perturbation method for motion of a spherical solid particle in plane couette fluid
   flow'. \M Jalaal , M Nejad , P Jalili , M Esmaeilpour , H Bararnia , E Ghasemi , S Soleimani , D Ganji , S
   Moghimi . Computers & Mathematics with Applications 2011. p. .
- [Kim and Karrila ()] 'Microhydrodynamics: principles and selected applications'. S Kim , S J Karrila . Bubbles,
   drops and particles in non-Newtonian fluids, 2007. 85 p. 251. (CANADIAN JOURNAL OF CHEMICAL
   ENGINEERING)
- 191 [Oseen ()] Neuere methoden und ergebnisse in der hydrodynamik, C W Oseen . 1927. Akad. Verlag-Ges.
- [Yaghoobi and Torabi ()] 'Novel solution for acceleration motion of a vertically falling nonspherical particle by
   VIM-Padé approximant'. H Yaghoobi , M Torabi . *Powder Technology* 2012. 215 p. .
- [Torabi and Yaghoobi ()] 'Novel solution for acceleration motion of a vertically falling spherical particle by HPM Padé approximant'. M Torabi , H Yaghoobi . Advanced Powder Technology 2011. 22 p. .
- [Stokes ()] On the effect of the internal friction of fluids on the motion of pendulums, G G Stokes . Press1851.
   Pitt.
- IJalaal and Ganji ()] 'On unsteady rolling motion of spheres in inclined tubes filled with incompressible
   Newtonian fluids'. M Jalaal , D D Ganji . Advanced Powder Technology 2011. 22 p. .
- 200 [Chien ()] 'Settling velocity of irregularly shaped particles'. S F Chien . SPE Drilling & Completion 1994. 9 p. .
- [Lovalenti and Brady ()] 'The hydrodynamic force on a rigid particle undergoing arbitrary timedependent motion at small Reynolds number'. P M Lovalenti , J F Brady . *Journal of Fluid Mechanics* 1993. 256 p. .
- [Sano ()] 'Unsteady flow past a sphere at low Reynolds number'. T Sano . Journal of Fluid Mechanics 1981. 112
   p. .